CLAIMS

We claim:

1. (canceled)	
2. (canceled)	
3. (canceled)	
4. (canceled)	
5. (canceled)	
6. (canceled)	
7. (canceled)	
8. (canceled)	
9. (canceled)	
10. (canceled)	
11. (canceled)	
12. (canceled)	
13. (canceled)	
14. (canceled)	
15. (canceled)	
16. (canceled)	
17. (canceled)	
18. (canceled)	
19. (canceled)	
20. (canceled)	
21. (canceled)	
	2

- 22. (canceled)
- 23. (canceled)
- 24. (canceled)
- 25. (canceled)
- 26. (canceled)
- 27. (canceled)
- 28. (canceled)
- 29. (canceled)
- 30. (canceled)
- 31. (canceled)
- 32. (canceled)
- 33. (canceled)
- 34. (canceled)
- 35. (canceled)
- 36. (canceled)
- 37. (canceled)
- 38. (canceled)
- 39. (canceled)
- 40. (canceled)
- 41. (canceled)
- 42. (canceled)
- 43. (canceled)

44. (original) A method for treating a patient for a bone related condition comprising the steps of: measuring a bone characteristic level in a bone of said patient to yield a T-score having a value;

if the T-score is abnormal, conducting a gait analysis to yield a gait characterization;

if the gait characterization is abnormal, measuring a bone marker concentration in at least one body fluid of said patient to yield a bone marker level having a value;

prescribing a therapy based on at least one of the said gait characterization, said T-score, and bone marker level; and

designating a future time to repeat said measurement of a bone characteristic level, said gait analysis, and said measurement of a bone marker concentration.

- 45. (original) The method of claim 44 wherein said future time to repeat said measurement of a bone characteristic level is during the twelfth month from the previous measurement.
- 46. (original) The method of claim 44 wherein the step of designating a future time to repeat said gait analysis includes scheduling a series of eight gait analyses over a period of time.
- 47. (original) The method of claim 44 wherein said future time to repeat said bone marker measurement is during the third month from the previous measurement.
- 48. (previously presented) The method of claim 44 wherein the bone characteristic level is measured using a bone characteristic measuring unit, comprising: a space for housing said bone of said patient; a positioning device for holding said bone; a plurality of ultrasound transducers for transmitting and detecting signals; and an output for outputting said first score value.
- 49. (previously presented) The method of claim 48 wherein the bone characteristic is a quantitative ultrasound index.
- 50. (previously presented) The method of claim 48 wherein the bone characteristic is a stiffness index.
- 51. (previously presented) The method of claim 44 wherein the bone characteristic level is measured using X-ray absorptiometry.
- 52. (previously presented) The method of claim 44 wherein the bone characteristic level is measured using quantitative ultrasonometry.
- 53. (previously presented) The method of claim 44 wherein the bone characteristic level is measured using quantitative computed tomography.
- 54. (previously presented) The method of claim 44 wherein the bone characteristic is bone mineral density.

- 55. (previously presented) The method of claim 44 further comprising the step of assessing a plurality of risk factors attributable to the patient.
- 56. (previously presented) The method of claim 55 wherein said therapy is prescribed based at least in part upon an assessment of patient risk factors.
- 57. (previously presented) The method of claim 44 wherein said therapy is prescribed based upon an output of an integrated unit having received the T-score, the gait characterization, and the bone marker level.
- 58. (previously presented) The method of claim 57, wherein said integrated unit comprises a receiver in data communication with a processing unit and a display unit in data communication with the processing unit.
- 59. (previously presented) The method of claim 44 wherein the bone marker concentration is measured by a bone marker measurement device, wherein said device comprises: a container containing a body fluid; a mechanism for holding the said container; an analyzer for determining a concentration of an absorbing constituent in a solution; and an output for outputting the bone marker level value.
- 60. (previously presented) The method of claim 44 wherein the gait analysis is characterized by a gait analysis procedure conducted on said patient having a balance, wherein said procedure comprises the steps of: examining the balance of the patient wherein the patient is standing on both feet; examining the balance of the patient wherein the patient is standing on a first foot; and examining the balance of the patient wherein the patient is standing on a second foot.
- 61. (previously presented) The method of claim 44 wherein the gait analysis is characterized by a gait analysis procedure conducted on said patient having a balance, wherein said procedure comprises the steps of: having the patient stand on a plurality of platforms; detecting pressure exerted on said plurality of platforms; and determining a pressure differential on said plurality of platforms.